



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> :  
 C07D 473/02, 211/26, 211/28  
 C07D 211/30, 277/20, 277/44  
 C07D 277/02, C07C 271/02, 271/08  
 C07C 271/10, 271/12, 271/14  
 C07C 271/18, 229/02, 229/24  
 C07C 229/26, 229/34, 229/40  
 C07C 233/04, 233/05, 233/06  
 C07C 233/07, 233/12, 233/31  
 C07C 233/47, 233/51, 235/74  
 C07C 235/76, 237/06, 237/08, 239/00

A1

(11) International Publication Number: WO 93/07148


(43) International Publication Date: 15 April 1993 (15.04.93)

(21) International Application Number: PCT/US92/08454

(22) International Filing Date: 5 October 1992 (05.10.92)

(30) Priority data:  
771,760 4 October 1991 (04.10.91) US(71) Applicants: SLOAN-KETTERING INSTITUTE FOR  
CANCER RESEARCH [US/US]; 1275 York Avenue,  
New York, NY 10021 (US). THE TRUSTEES OF CO-  
LUMBIA UNIVERSITY IN THE CITY OF NEW  
YORK [US/US]; West 116th Street & Broadway, New  
York, NY 10027 (US).(72) Inventors: BRESLOW, Ronald ; 275 Broad Avenue, Eng-  
lewood, NJ 07631 (US). MARKS, Paul, A. ; Beach Hill  
Road, Bridgewater, CT 06752 (US). RIFKIND, Ri-  
chard, A. ; 30 Sutton Place, New York, NY 10022 (US).  
JURSIC, Branko ; 91 Spanish Fort Boulevard, New Or-  
leans, LA 70124 (US).(74) Agent: WHITE, John, P.; Cooper & Dunham, 30 Rocke-  
feller Plaza, New York, NY 10112 (US).(81) Designated States: AU, CA, FI, HU, JP, KR, NO, RU, Eu-  
ropean patent (AT, BE, CH, DE, DK, ES, FR, GB, GR,  
IE, IT, LU, MC, NL, SE).**Published***With international search report.**Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.*

(54) Title: NOVEL POTENT INDUCERS OF TERMINAL DIFFERENTIATION AND METHODS OF USE THEREOF



## (57) Abstract

The present invention provides the compound having structure (I), wherein each of R<sub>1</sub> and R<sub>2</sub> are independently the same as or different from each other; when R<sub>1</sub> and R<sub>2</sub> are the same, each is a substituted or unsubstituted arylamino, cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; when R<sub>1</sub> and R<sub>2</sub> are different, R<sub>1</sub> = R<sub>3</sub>-N-R<sub>4</sub>, wherein each of R<sub>3</sub> and R<sub>4</sub> are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or pyridine group, or R<sub>3</sub> and R<sub>4</sub> bond together to form a piperidine group and R<sub>2</sub> is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and n is an integer from about 4 to about 8. The present invention also provides a method of selectively inducing terminal differentiation of neoplastic cells and thereby inhibiting proliferation of such cells. Moreover, the present invention provides a method of treating a patient having a tumor characterized by proliferation of neoplastic cells. Lastly, the present invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically acceptable amount of the compound above.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SK	Slovak Republic
CI	Côte d'Ivoire	LK	Sri Lanka	SN	Senegal
CM	Cameroon	LU	Luxembourg	SU	Soviet Union
CS	Czechoslovakia	MC	Monaco	TD	Chad
CZ	Czech Republic	MG	Madagascar	TG	Togo
DE	Germany	ML	Mali	UA	Ukraine
DK	Denmark	MN	Mongolia	US	United States of America
ES	Spain			VN	Viet Nam
FI	Finland				

NOVEL POTENT INDUCERS OF TERMINAL DIFFERENTIATION AND  
METHODS OF USE THEREOF

Background of the Invention

5

Throughout this application various publications are referenced by arabic numerals within parentheses. Full citations for these publications may be found at the end of the specification immediately preceding the claims.

10 The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

15 Cancer is a disorder in which a population of cells has become, in varying degrees, unresponsive to the control mechanisms which normally govern proliferation and differentiation. For many years there have been two principal strategies for chemotherapeutic treatment of  
20 cancer: a) blocking hormone-dependent tumor cell proliferation by interference with the production or peripheral action of sex hormones; and b) killing cancer cells directly by exposing them to cytotoxic substances, which injure both neoplastic and normal cell populations.

25

Relatively recently, cancer therapy is also being attempted by the induction of terminal differentiation of the neoplastic cells (1). In cell culture models  
30 differentiation has been reported by exposure of cells to a variety of stimuli, including: cyclic AMP and retinoic acid (2,3), aclarubicin and other anthracyclines (4).

There is abundant evidence that neoplastic transformation  
35 does not necessarily destroy the potential of cancer cells to differentiate (1,5,6). There are many examples of tumor cells which do not respond to the normal

**SUBSTITUTE SHEET**

regulators of proliferation and appear to be blocked in the expression of their differentiation program, and yet can be induced to differentiate and cease replicating. A variety of agents, including some relatively simple polar compounds (5,7-9), derivatives of vitamin D and retinoic acid (10-12), steroid hormones (13), growth factors (6,14), proteases (15,16), tumor promoters (17,18), and inhibitors of DNA or RNA synthesis (4,19-24), can induce various transformed cell lines and primary human tumor explants to express more differentiated characteristics.

Early studies by the present inventors identified a series of polar compounds that were effective inducers of differentiation in a number of transformed cell lines (8,9). Of these, the most effective induce, was the hybrid polar/apolar compound N,N'-hexamethylene bsacetamide (HMBA) (9). The use of this polar/apolar compound to induce murine erythroleukemia cells (MELC) to undergo erythroid differentiation with suppression of oncogenicity has proved a useful model to study inducer-mediated differentiation of transformed cells (5,7-9). HMBA-induced MELC terminal erythroid differentiation is a multistep process. Upon addition of HMBA to MELC (745A-DS19) in culture, there is a latent period of 10 to 12 hours before commitment to terminal differentiation is detected. Commitment is defined as the capacity of cells to express terminal differentiation despite removal of inducer (25). Upon continued exposure to HMBA there is progressive recruitment of cells to differentiate. The present inventors have reported that MELC cell lines made resistant to relatively low levels of vincristine become markedly more sensitive to the inducing action of HMBA and can be induced to differentiate with little or no latent period (26).

HMBA is capable of inducing phenotypic changes consistent

**SUBSTITUTE SHEET**

with differentiation in a broad variety of cells lines (5). The characteristics of the drug induced effect have been most extensively studied in the murine erythroleukemia cell system (MELC) (5,25,27,28). MELC  
5 induction of differentiation is both time and concentration dependent. The minimum concentration required to demonstrate an effect in vitro in most strains is 2 to 3 mM; the minimum duration of continuous exposure generally required to induce differentiation in  
10 a substantial portion (>20%) of the population without continuing drug exposure is about 36 hours.

The primary target of action of HMBA is not known. There is evidence that protein kinase C is involved in the  
15 pathway of inducer-mediated differentiation (29). The in vitro studies provided a basis for evaluating the potential of HMBA as a cytodifferentiation agent in the treatment of human cancers (30). Several phase I clinical trials with HMBA have been completed (31-36).  
20 Clinical trials have shown that this compound can induce a therapeutic response in patients with cancer (35,36). However, these phase I clinical trials also have demonstrated that the potential efficacy of HMBA is limited, in part, by dose-related toxicity which prevents  
25 achieving optimal blood levels and by the need for intravenous administration of large quantities of the agent, over prolonged periods.

Recently, the present inventors have reported a number of  
30 compounds related to HMBA with polar groups separated by apolar linkages that, on a molar basis, are as active (37) or 100 times more active than HMBA (38). As a class, however, it has been found that the symmetrical dimers such as HMBA and related compounds are not the  
35 best cytodifferentiating agents.

It has unexpectedly been found that the best compounds

**SUBSTITUTE SHEET**



-4-


comprise two polar end groups separated by a flexible chain of methylene groups, wherein one or both of the polar end groups is a large hydrophobic group. Preferably, the polar end groups are different and only one is a large hydrophobic group. These compounds are unexpectedly a thousand times more active than HMBA and ten times more active than HMBA related compounds.

This new class of compounds of the present invention may be useful for selectively inducing terminal differentiation of neoplastic cells and therefore aid in treatment of tumors in patients.

**SUBSTITUTE SHEET**

Summary of the Invention

The present invention provides the compound having the structure:




10

herein each of  $\text{R}_1$  and  $\text{R}_2$  are independently the same as or different from each other; when  $\text{R}_1$  and  $\text{R}_2$  are the same, each is a substituted or unsubstituted arylamino, cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; when  $\text{R}_1$  and  $\text{R}_2$  are different,  $\text{R}_1 = \text{R}_3\text{-N-R}_4$ , wherein each of  $\text{R}_3$  and  $\text{R}_4$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or pyridine group, or  $\text{R}_3$  and  $\text{R}_4$  bond together to form a piperidine group and  $\text{R}_2$  is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and  $n$  is an integer from about 4 to about 8.

25

The present invention also provides the compound above having the structure:



30


wherein each of  $\text{R}_3$  and  $\text{R}_4$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched

-6-

or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or pyridine group, or  $R_3$  and  $R_4$  bond together to form a piperidine group;  $R_2$  is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and  $n$  is an integer from about 4 to about 8.

The present invention also provides the compound above having the structure:

10




15

wherein  $R$  is a substituted or unsubstituted arylamino, cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; and  $n$  is an integer from about 4 to about 8.

20

The present invention also provides the compound having the structure:




wherein each of  $X$  and  $Y$  are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group;  $R$  is a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of  $m$  and  $n$  are independently the same as or

**SUBSTITUTE SHEET**



different from each other and are each an integer from about 0 to about 8.


The present invention further provides the compound  
5 having the structure:



10

wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of R<sub>1</sub> and R<sub>2</sub> are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m, n, and o are independently the same as or different from each other and are each an integer from about 0 to about 8.

The present invention still further provides the compound  
25 having the structure:




30

wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino,

alkyloxyalkylamino, or aryloxyalkylamino group; each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or  
 5 aryloxy group; and each of  $m$  and  $n$  are independently the same as or different from each other and are each an integer from about 0 to about 8.

The present invention also provides the compound having  
 10 the structure:




15

wherein each of  $X$  and  $Y$  are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino,  
 20 alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; and each of  $m$  and  $n$  are independently the same as or different from each other and are each an integer from about 0 to about 8.

25

The present invention also provides the compound having the structure:




30

wherein each of  $X$  and  $Y$  are independently the same as or different from each other and are a hydroxyl, amino or  
 35 hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino,

alkyloxyalkylamino, or aryloxyalkylamino group; each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of  $m$  and  $n$  are independently the same as or different from each other and are each an integer from about 0 to about 8.


The present invention further provides the compound having the structure:



15

wherein each of  $X$  and  $Y$  are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamine, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; and  $n$  is an integer from about 0 to about 8.

The present invention still further provides the compound having the structure:




30

wherein each of  $X$  and  $Y$  are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamine, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of

-10-

$R_1$  and  $R_2$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, aryloxy, carbonylhydroxylamino, or fluoro group; and  
5 each of  $m$  and  $n$  are independently the same as or different from each other and are each an integer from about 0 to about 8.

10 The present invention also provides the compound having the structure:




15

wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino,  
20 arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

The present invention also provides the compound having the structure:

25




30

wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino,  
35 alkyloxyalkylamino, or aryloxyalkylamino group.

The present invention further provides the compound

**SUBSTITUTE SHEET**

having the structure:



wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

10

In addition, the present invention provides a method of selectively inducing terminal differentiation of neoplastic cells and thereby inhibiting proliferation of such cells which comprises contacting the cells under suitable conditions with an effective amount of any of the compounds above, effective to selectively induce terminal differentiation.

15

20

The present invention also provides a method of treating a patient having a tumor characterized by proliferation of neoplastic cells which comprises administering to the patient an effective amount of any of the compounds above, effective to selectively induce terminal differentiation of such neoplastic cells and thereby inhibit their proliferation.

25


Lastly, the present invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically acceptable amount of any of the compounds above.

30

Detailed Description of the Invention

The present invention provides the compound having the structure:


5



- 10 wherein each of  $\text{R}_1$  and  $\text{R}_2$  are independently the same as or different from each other; when  $\text{R}_1$  and  $\text{R}_2$  are the same, each is a substituted or unsubstituted arylamino, cycloalkyl-amino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; when  $\text{R}_1$  and  $\text{R}_2$  are
- 15 different,  $\text{R}_1 = \text{R}_3\text{-N-R}_4$ , wherein each of  $\text{R}_3$  and  $\text{R}_4$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or
- 20 pyridino group, or  $\text{R}_3$  and  $\text{R}_4$  bond together to form a piperidine group and  $\text{R}_2$  is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and  $n$  is an integer from about 4 to about 8.

- 25 The present invention also provides the compound above having the structure:

30



- wherein each of  $\text{R}_3$  and  $\text{R}_4$  are independently the same as or different from each other and are a hydrogen atom, a
- 35 hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or pyridine group, or  $\text{R}_3$  and  $\text{R}_4$

**SUBSTITUTE SHEET**



bond together to form a piperidine group;  $R_2$  is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and  $n$  is an integer from about 4 to about 8.

5

In the preferred embodiment of the compound above,  $R_2$  is a hydroxylamino, hydroxyl, amino, methylamino, dimethylamino, or methoxy group and  $n$  is 6. Most preferably,  $R_4$  is a hydrogen atom and  $R_3$  is a substituted or unsubstituted phenyl group.

10

The phenyl group may be substituted with a methyl, cyano, nitro, trifluoromethyl, amino, aminocarbonyl, methylcyano, chloro, fluoro, bromo, iodo, 2,3-difluoro, 2,4-difluoro, 2,5-difluoro, 3,4-difluoro, 3,5-difluoro, 2,6-difluoro, 1,2,3-trifluoro, 2,3,6-trifluoro, 2,4,6-trifluoro, 3,4,5-trifluoro, 2,3,5,6-tetrafluoro, 2,3,4,5,6-pentafluoro, azido, hexyl, t-butyl, phenyl, carboxyl, hydroxyl, methoxy, benzyloxy, phenylaminooxy, phenylmethoxy, phenylamino-carbonyl, methoxycarbonyl, methylaminocarbonyl, dimethylamino, dimethylaminocarbonyl, or hydroxylamino-carbonyl group.

15

20

In other preferred embodiments of the compound above,  $R_4$  is a hydrogen atom and  $R_3$  is a cyclohexyl group;  $R_4$  is a hydrogen atom and  $R_3$  is a methoxy group;  $R_3$  and  $R_4$  each bond together to form a piperidine group;  $R_4$  is a hydrogen atom and  $R_3$  is a hydroxyl group;  $R_4$  is a hydrogen atom and  $R_3$  is a benzyloxy group;  $R_4$  is a hydrogen atom and  $R_3$  is a  $\delta$ -pyridine group;  $R_4$  is a hydrogen atom and  $R_3$  is a  $\beta$ -pyridine group;  $R_4$  is a hydrogen atom and  $R_3$  is a  $\alpha$ -pyridine group;  $R_3$  and  $R_4$  are both methyl groups; or  $R_4$  is a methyl group and  $R_3$  is a phenyl group.


25

30

-14-

The present invention also provides the compound having the structure:

5




10 wherein R is a substituted or unsubstituted arylamino, cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; and n is an integer from about 4 to about 8.

15 In the preferred embodiment of the compound above, R is a substituted or unsubstituted phenylamino group. The phenylamino group may be substituted with a cyano, methylcyano, nitro, carboxyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, trifluoromethyl, hydroxylaminocarbonyl, N-hydroxylaminocarbonyl, methoxycarbonyl, chloro, fluoro, methyl, methoxy, 2,3-difluoro, 2,3-difluoro, 2,4-difluoro, 2,5-difluoro, 2,6-difluoro, 3,5-difluoro, 2,6-difluoro, 2,3,6-trifluoro, 1,2,3-trifluoro, 3,4,5-trifluoro, 2,3,4,5-tetrafluoro, or  
25 2,3,4,5,6-pentafluoro group.

In another embodiment of the compound above, R is a cyclohexylamino group.

30 The present invention also provides the compound having the structure:




**SUBSTITUTE SHEET**

wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; R is a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

In the preferred embodiment of the compound above, each of X, Y, and R is a hydroxyl group and each of m and n is 5.

The present invention also provides the compound having the structure:




wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of R<sub>1</sub> and R<sub>2</sub> are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m, n, and o are independently the same as or different from each other and are each an integer from about 0 to about 8.

In the preferred embodiment of the compound above, each of X and Y is a hydroxyl group and each of R<sub>1</sub> and R<sub>2</sub> is a methyl group. Most preferably, each of n and o is 6, and m is 2.

5

The present invention also provides the compound having the structure:

10




15 wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of R<sub>1</sub> and R<sub>2</sub> are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

25

The present invention also provides the compound having the structure:

30



35


wherein each of X and Y are independently the same as or

different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

In the preferred embodiment of the compound above, each of X and Y is a hydroxyl group and each of m and n is 5.

The present invention also provides the compound having the structure:


15



wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of R<sub>1</sub> and R<sub>2</sub> are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

The present invention also provides the compound having the structure:

5




10 wherein each of X and Y are independently the same as or  
different from each other and are a hydroxyl, amino or  
hydroxylamino group, a substituted or unsubstituted  
alkyloxy, alkylamino, dialkylamino, arylamino,  
alkylaryl amino, alkyloxyamino, aryloxyamino,  
15 alkyloxyalkylamino, or aryloxyalkylamino group; and n is  
an integer from about 0 to about 8.

In the preferred embodiment of the compound above, each of X and Y is a dimethylamino group and n is 4 or 5.

20

The present invention also provides the compound having the structure:



wherein each of X and Y are independently the same as or  
30 different from each other and are a hydroxyl, amino or  
hydroxylamino group, a substituted or unsubstituted  
alkyloxy, alkylamino, dialkylamino, arylamino,  
alkylaryl amino, alkyloxyamino, aryloxyamino,  
alkyloxyalkylamino, or aryloxyalkylamino group; each of  
35  $R_1$  and  $R_2$  are independently the same as or different from  
each other and are a hydrogen atom, a hydroxyl group, a  
substituted or unsubstituted alkyl, aryl, alkyloxy,

# SUBSTITUTE SHEET



aryloxy, carbonylhydroxylamino, or fluoro group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.


5

In the preferred embodiment of the compound above, each of X and Y is a hydroxylamino group,  $R_1$  is a methyl group,  $R_2$  is a hydrogen atom, and each of m and n is 2. In another preferred embodiment, each of X and Y is a hydroxylamino group,  $R_1$  is a carbonylhydroxylamino group,  $R_2$  is a hydrogen atom, and each of m and n is 5. In a further preferred embodiment, each of X and Y is a hydroxylamino group, each of  $R_1$  and  $R_2$  is a fluoro group, and each of m and n is 2.

15

The present invention also provides the compound having the structure:

20




wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

30

Preferably,  $R_1$  is a phenylamino group and  $R_2$  is a hydroxylamino group.

The present invention also provides the compound having the structure:




10 wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

15

Preferably,  $R_1$  is phenylamino group and  $R_2$  is hydroxylamino group.

20 The present invention also provides the compound having the structure:



30 wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

35 In the preferred embodiment, either  $R_1$  or  $R_2$  is a hydroxylamino group.

The present invention also provides a method of

selectively inducing terminal differentiation of neoplastic cells and thereby inhibiting proliferation of such cells which comprises contacting the cells under suitable conditions with an effective amount of any of the compounds above, effective to selectively induce terminal differentiation.

The contacting must be performed continuously for a prolonged period of time, i.e. for at least 48 hours, preferably for about 4-5 days or longer.

The method may be practiced in vivo or in vitro. If the method is practiced in vitro, contacting may be effected by incubating the cells with the compound. The concentration of the compound in contact with the cells should be from about 1  $\mu$ M to about 25 mM, preferably from 4  $\mu$ M to about 5 mM. The concentration depends upon the individual compound and the state of the neoplastic cells.

The method may also comprise initially treating the cells with an antitumor agent so as to render them resistant to an antitumor agent and subsequently contacting the resulting resistant cells under suitable conditions with an effective amount of any of the compounds above, effective to selectively induce terminal differentiation of such cells.

The antitumor agent may be one of numerous chemotherapy agents such as an alkylating agent, an antimetabolite, a hormonal agent, an antibiotic, colchicine, a vinca alkaloid, L-asparaginase, procarbazine, hydroxyurea, mitotane, nitrosoureas or an imidazole carboxamide. Suitable agents are those agents which promote depolarization of tubulin. Preferably the antitumor agent is colchicine or a vinca alkaloid; especially preferred are vinblastine and vincristine. In

embodiments where the antitumor agent is vincristine, the cells preferably are treated so that they are resistant to vincristine at a concentration of about 5 mg/ml. The treating of the cells to render them resistant to an antitumor agent may be effected by contacting the cells with the agent for a period of at least 3-5 days. The contacting of the resulting cells with any of the compounds above is performed as described previously.

10 The present invention also provides a method of treating a patient having a tumor characterized by proliferation of neoplastic cells which comprises administering to the patient an effective amount of any of the compounds above, effective to selectively induce terminal  
15 differentiation of such neoplastic cells and thereby inhibit their proliferation.

The method of the present invention is intended for the treatment of human patients with tumors. However, it is also likely that the method would be effective in the treatment of tumors in other mammals. The term tumor is intended to include any cancer caused by the proliferation of neoplastic cells, such as lung cancer, acute lymphoid myeloma, bladder melanoma, renal  
20 carcinoma, breast carcinoma, or colorectal carcinoma. The administration of the compound to the patient may be effected orally or parenterally. To date, administration intravenously has proven to be effective. The administration of the compound must be performed  
25 continuously for a prolonged period of time, such as for at least 3 days and preferably more than 5 days. In the most preferred embodiments, the administration is effected continuously for at least 10 days and is repeated at intervals wherein at each interval the  
30 administration is continuously effected for at least 10 days. For example, the administration may be effected at intervals as short as 5-10 days, up to about 25-35 days

and continuously for at least 10 days during each such interval. The optimal interval period will vary depending on the type of patient and tumor. For example, in the incidence of acute leukemia, the so called  
5 myelodysplastic syndrome, continuous infusion would seem to be indicated so long as the patient tolerated the drug without toxicity and there was a positive response.

The amount of the compound administered to the patient is  
10 less than an amount which would cause toxicity in the patient. In the certain embodiments, the amount of the compound which is administered to the patient is less than the amount which causes a concentration of the compound in the patient's plasma to equal or exceed the  
15 toxic level of the compound. Preferably, the concentration of the compound in the patient's plasma is maintained at about 1.0 mM. It has been found with HMBA that administration of the compound in an amount from  
20 about 5 gm/m<sup>2</sup>/day to about 30 gm/m<sup>2</sup>/day, particularly about 20 gm/m<sup>2</sup>/day, is effective without producing toxicity in the patient. The optimal amount of the compound which should be administered to the patient in the practice of the present invention will depend on the particular compound used and the type of cancer being  
25 treated.

This invention, in addition to the above listed compounds, is intended to encompass the use of homologs and analogs of such compounds. In this context, homologs  
30 are molecules having substantial structural similarities to the above-described compounds and analogs are molecules having substantial biological similarities regardless of structural similarities.

35 The method may also comprise initially administering to the patient an amount of an antitumor agent to render the cells resistant to an antitumor agent and subsequently

**SUBSTITUTE SHEET**



administering to the patient an effective amount of any of the compounds above, effective to selectively induce terminal differentiation of such neoplastic cells and thereby inhibit their proliferation.

5

The antitumor agent may be one of numerous chemotherapy agents such as an alkylating agent, an antimetabolite, a hormonal agent, an antibiotic, colchicine, a vinca alkaloid, L-asparaginase, procarbazine, hydroxyurea, mitotane, nitrosoureas or an imidazole carboxamide. Suitable agents are those agents which promote depolarization of tubulin. Preferably the antitumor agent is colchicine or a vinca alkaloid; especially preferred are vinblastine and vincristine. In  
10  
15  
20  
embodiments where the antitumor agent is vincristine, an amount is administered to render the cells resistant to vincristine at a concentration of about 5 mg/ml. The administration of the agent is performed essentially as described above for the administration of any of the compounds. Preferably, the administration of the agent is for a period of at least 3-5 days. The administration of any of the compounds above is performed as described previously.

25 The present invention also provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier, such as sterile pyrogen-free water, and a therapeutically acceptable amount of any of the compounds above. Preferably, the effective amount is an amount  
30  
effective to selectively induce terminal differentiation of suitable neoplastic cells and less than an amount which causes toxicity in a patient.

35 Lastly, the present invention provides the pharmaceutical composition above in combination with an antitumor agent. The antitumor agent may be any of the agents previously described.

**SUBSTITUTE SHEET**



The invention is illustrated in the Experimental Details section which follows. This section is set forth to aid in an understanding of the invention but is not intended to, and should not be construed to, limit in any way the invention as set forth in the claims which follow thereafter.

Experimental DetailsCells and Materials

5 MELC 745A-DS19 cells and the variants of MELC derived  
from this cell line, namely, the vincristine-resistant  
MELC V3.17 and VCR.C(2)15 cell lines (26), and the  
dimethylsulfoxide-resistant cell line, DR10 (39), were  
maintained in alpha minimal essential medium containing  
10 10% fetal calf serum (16). Cell cultures for all  
experiments were initiated with cells in logarithmic  
growth phase (day 2 cultured cells) at a density of  $10^5$   
cells/ml. Inducer compounds were added in the final  
concentrations indicated below, dissolved in culture  
15 medium without fetal calf serum unless otherwise  
indicated. Cell density and benzidine reactivity were  
determined as described (16).


Commitment to terminal differentiation, characterized by  
20 limited cell division (colony size <32 cells) and  
accumulation of hemoglobin (benzidine reactive colonies)  
was assayed by a colony cloning assay using 2%  
methylcellulose as described (25) (see Table 1 for  
results).

25 HL-60 human leukemia cells, derived from peripheral blood  
leukocytes of a patient with acute promyelocytic leukemia  
(40). Induced differentiation of HL-60 cells assayed by  
determining the proportion of cells that developed the  
30 capacity to reduce nitroblue tetrazolium (NBT) (41) (see  
Table 2 for results).

Chemistry


The compounds having the structure:

35



Preparation of  $\text{PhCH}_2\text{ONHOC}(\text{CH}_2)_6\text{COOCH}_3$ :

A solution of suberic acid monomethyl ester (1.9 g; 0.01 mol), oxaloyl chloride (1.75 mL; 2.54 g; 0.02 mol) and 0.1 mL DMF in benzene (200 mL) was stirred overnight at room temperature. The solvent was evaporated and oily residue was dissolved in chloroform (~20 mL) and mixed together with chloroform solution (100 mL) of O-benzylhydroxylamine (2.46 g; 0.02 mol) and pyridine (1.6 mL; 1.68 g; 0.02 mol). The reaction mixture was stirred at room temperature overnight. The chloroform solution was washed with water (50 mL), 10% hydrochloric acid, and again with water (2 x 50 mL). The organic layer was dried over anhydrous magnesium sulfate and evaporated. The solid residue was slurried in hexanes (~100 mL) and filtered. The yield of  $\text{PhCH}_2\text{ONHOC}(\text{CH}_2)_6\text{COOCH}_3$  was 2.61 g (89%).




The above suberic acid monobenzyloxyamide monomethyl ester (1 g; 3.4 mol) was dissolved in dry methanol (50 mL) and 5% Pd-C (50 mg) was added. The black suspension was shaken under hydrogen pressure (~50 psi) overnight at room temperature. The catalyst was separated by filtration, and filtrate was evaporated. The solid residue was slurried in hexanes (~20 mL) and filtered. The yield of the monomethyl ester monohydroxamic acid of suberic acid was 900 mg (95%).

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.31 (s, NHOH, 1H);

-28-

8.89 (s, broad, NHOH, 1H); 3.57 (s, CH<sub>3</sub>, 3H); 2.27 (t, J=7.4Hz, CH<sub>2</sub>COOCH<sub>3</sub>, 2H); 1.91 (t, J=7.4Hz, CH<sub>2</sub>CONHOH, 2H); 1.49 (m, 4H), 1.24 (m, 4H).

5




10

Suberic acid monobenzyloxyamide monomethyl ester (1g; 3.4 mmol) and potassium hydroxide (210 mg; 3.75 mmol) were dissolved in 10 mL of methanol-water (4:1) mixture. The reaction mixture was refluxed two hours and solvent was evaporated. The solid residue was dissolved in 5 mL water and acidified with conc. hydrochloric acid to pH 5. White precipitate was filtered, dried and crystallized from ethyl acetate-hexanes. The yield of suberic acid monobenzyloxyamide was 820 mg (86%). The product was dissolved in methanol (50 mL) and 5% Pd-C (50 mg) was added. The reaction mixture was shaken under hydrogen pressure (50 psi) overnight. The catalyst was separated by filtration and filtrate was evaporated. The solid residue was slurried in hexanes and filtered. The yield of suberic acid monohydroxamic acid was 520 mg (81%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 11.96 (s, broad, COOH, 1H); 10.31 (s, NHOH, 1H); 8.63 (s, broad, NHOH, 1H); 2.17 (s, J=7.4Hz, CH<sub>2</sub>COOH, 2H); 1.91 (s, CH<sub>2</sub>CONHOH, 2H); 1.46 (m, 4H); 1.22 (m, 4H).

30

Compounds having the structure:


35




**SUBSTITUTE SHEET**

General Procedure

A pyridine (500 mL) solution of O-benzylhydroxylamine (2.46 g; 0.02 mol), the corresponding amine (0.02 mol) and suberoyl chloride was stirred at room temperature overnight. The solvent was evaporated and the semisolid residue was dissolved in 1000 mL chloroform-methanol (4:1); the resulting solution was washed with water (2 x 100 mL), 10% hydrochloric acid (3 x 100 mL), and again with water (2 x 100 mL). Organic layer was dried over anhydrous magnesium sulfate and evaporated. The solid residue was dissolved in methanol (100 mL) and 5% Pd-C was added. The black suspension was shaken under hydrogen pressure (~50 psi) overnight. The catalyst was separated by filtration, and the filtrate was evaporated. The target products were isolated by column chromatography on silica gel with ethyl acetate-tetrahydrofuran.




Yield 1.1 g (26%).  $^1\text{H}$  NMR (DMSO- $\text{D}_6$ , 200 MHz),  $\delta$ (ppm) 10.93 (s,  $\text{NHOCH}_3$ , 1H); 10.32 (s,  $\text{NHOH}$ , 1H); 8.66 (s,  $\text{NHOH}$ , 1H); 3.55 (s,  $\text{CH}_3$ , 3H); 1.91 (t,  $J=7.6\text{Hz}$ ,  $\text{CH}_2\text{CO-}$ , 4H); 1.45 (m, 4H); 1.20 (m, 4H).


**SUBSTITUTE SHEET**

-30-

Yield 1.2 g (21%).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 200 MHz),  $\delta$ (ppm)  
 10.31 (s, NHOH, 1H); 8.60 (s, broad, NHOH, 1H); 7.57 (d,  
 $J=7.6\text{Hz}$ ,  $\text{NH}-\text{C}_6\text{H}_{11}$ , 1H), 3.40 (m,  $\text{CH}-\text{NH}$ , 1H); 1.99 (t,  
 $J=7\text{Hz}$ ,  $\text{CH}_2\text{CONHC}_6\text{H}_{11}$ , 2H); 1.91 (t,  $J=7.6\text{Hz}$ ,  $\text{CH}_2\text{CONHOH}$ , 2H);  
 5 1.63 (m, 4H); 1.44 (m, 6H); 1.20 (m, 8H).




Yield 870 mg (20%).  $^1\text{H}$  NMR ( $\text{DMSO}-D_6$ , 200 MHz),  $\delta$ (ppm)  
 10.31 (s, NHOH, 1H); 8.67 (s, broad, NHOH, 1H); 2.85 (d,  
 15  $J=30\text{Hz}$ ,  $\text{N}(\text{CH}_3)_2$ , 6H); 2.24 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{CON}(\text{CH}_3)_2$ , 2H);  
 1.91 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{COONHOH}$ , 2H); 1.50 (m, 4H); 1.20 (m,  
 4H).



25 Yield 1.4 g (27%);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 200 MHz),  $\delta$ (ppm) 10.31  
 (s, NHOH, 1H); 8.67 (s, NHOH, 1H); 3.40 (2t,  $\text{CH}_2\text{N}$ , 4H);  
 2.20 (t,  $J=7.4\text{ Hz}$ ,  $\text{CH}_2\text{CON}(\text{CH}_2)_5$ , 2H); 1.91 (t,  $J=7.4\text{Hz}$ ,  
 $\text{CH}_2\text{CONHOH}$ , 2H); 1.10-1.60 (m, broad, 14 H).

30 Compound having structure:




**SUBSTITUTE SHEET**



-31-

The chloroform (500 mL) solution of O-benzylhydroxylamine (1.23 g; 0.01 mol), O-(trimethylsilyl)hydroxylamine (1.1 g; 0.01 mol), pyridine (1.6 mL; 1.7 g; 0.02 mol) and suberoyl chloride (1.8 mL; 2.11 g; 0.01 mol) was stirred at room temperature overnight. The reaction suspension was diluted with methanol (100 mL), washed with 10% hydrochloric acid (3 x 100 mL). The organic layer was dried over anhydrous magnesium sulfate and evaporated. The solid residue was subjected to chromatography on silica gel in ethyl acetate-tetrahydrofuran (4:1). The yield was 500 mg (17%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 11.09 (s, NHOCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, 1H); 10.31 (s, NHOH, 1H); 8.67 (s, broad, NHOH, 1H); 7.36 (s, C<sub>6</sub>H<sub>5</sub>, 5H), 4.76 (s, CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, 2H); 1.92 (t, J=7.4Hz, CH<sub>2</sub>CO-, 4H); 1.45 (m, 4H); 1.20 (m, 4H).

Compound having the structure:



25 Into a cooled solution of potassium hydroxide (2.24 g; 0.04 mol) and O-benzylhydroxylamine hydrochloride in 30 mL of tetrahydrofuran-water (1:1) mixture, 6-bromohexanoyl chloride (3.1 mL; 4.27 g; 0.02 mol) was added. The reaction mixture was stirred at room temperature for one hour. The solvent was evaporated and solid residue was partitioned between chloroform (200 mL) and water (100 mL). Chloroform layer was washed with 10% hydrochloric acid (3 x 50 mL) and water (2 x 50 mL). The organic layer was dried over anhydrous magnesium sulfate and evaporated. The product was purified by crystallization from ethyl acetate-hexanes. The yield of N-benzyl-6-bromohexanamide was 4.7 g (78%). A


**SUBSTITUTE SHEET**

-32-

dimethylsulfoxide (250 mL) solution of N-benzyloxy-6-bromohexanoyl amide (4.5 g; 15 mmol) and sodium cyanide (7.35 g; 0.15 mol) was heated at 130°C overnight. The solvent was evaporated and solid residue was partitioned  
5 between chloroform (300 mL) and water (300 mL). The chloroform layer was washed with water (5 x 100 mL), dried over anhydrous magnesium sulfate, and evaporated. The oily residue was purified by column chromatography on silica gel in ethyl acetate-tetrahydrofuran (4:1) as an  
10 eluent. The yield of N-benzyloxy-6-cyanohehexanoylamide was 1.62 g (43%). The product was dissolved in methanol (50 mL) and 5% Pd-C (100 mg) was added. The black suspension was shaken under hydrogen pressure (~50 psi) overnight. The catalyst was isolated by filtration and  
15 filtrate was evaporated. The solid residue was slurried in hexanes (~20 mL) and filtered. The yield of N-hydroxy-6-cyanohehexanoylamide was 900 mg (overall yield 30%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 10.32 (s, NHOH, 1H); 8.65 (s, NHOH, 1H); 2.45 (t, J=7Hz, CH<sub>2</sub>CN, 2H) 1.93  
20 (t, J=7Hz, CH<sub>2</sub>CONHOH, 2H); 1.49 (m, 4H); 1.33 (m, 2H).

Compounds having the structure:

25




30 General Procedure

A diacid dichloride (0.01 mol) was added into a cooled (0°C) solution of potassium hydroxide (1.12 g; 0.02 mol) and corresponding amine (0.01 mol) in 30 mL of  
35 tetrahydrofuran-water (1:1) mixture. The reaction mixture was stirred at room temperature about one hour. Solvent was evaporated and the solid residue was

**SUBSTITUTE SHEET**


partitioned between chloroform (300 mL) and water (300 mL). In some cases a small amount of methanol is necessary to dissolve all solid. The organic layer was washed with 10% potassium hydroxide (3 x 30 mL). The basic water extract was acidified with 10% hydrochloric acid. The precipitate was collected by filtration, dried and purified by crystallization from ethyl acetate or by column chromatography on silica gel in ethyl acetate-tetrahydrofuran (4:1). The yields are from 20-37%.

10




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 11.97 (s, COOH, 1H); 9.84 (s, NH, 1H); 7.57 (d, J=7.4Hz, ortho aromatic protons, 2H); 7.26 (t, J=8.4Hz, meta aromatic protons, 2H); 6.99 (t, J=7.4Hz, para aromatic proton, 1H), 2.27 (t, J=7Hz, CH<sub>2</sub>CONHPh, 2H); 2.18 (t, J=7.2Hz, 2H); 1.52 (m, 4H); 1.28 (m, 4H).

25




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 11.95 (s, COOH, 1H); 10.20 (s, NH, 1H); 8.10 (s, aromatic proton, 1H); 7.75 (m, aromatic proton, 1H); 7.45 (m, aromatic proton, 2H); 2.28 (t, J=7.4Hz, CH<sub>2</sub>CONHAr, 2H); 2.21 (t, J=7.2Hz, CH<sub>2</sub>COOH, 2H); 1.46 (m, 4H); 1.20 (m, 4H).

40



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 11.95 (s, COOH, 1H); 10.29 (s, NH, 1H); 7.75 (s, aromatic protons, 4H); 2.33 (t, J=7.2Hz, CH<sub>2</sub>CONHAr, 2H); 2.18 (t, J=7.4Hz, CH<sub>2</sub>COOH, 2H); 1.53 (m, 4H); 1.27 (m, 4H).


5



10


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200MHz), 11.98 (s, broad, COOH, 1H); 10.48 (s, NH, 1H); 8.21 (d, J=9.2Hz, aromatic protons, 2H); 7.82 (d, J=9.2Hz, aromatic proton, 2H); 2.36 (t, J=7.4Hz, CH<sub>2</sub>CONHAr, 2H); 2.18 (t, J=7.2Hz, CH<sub>2</sub>COOH, 2H); 1.55 (m, 4H); 1.29 (m, 4H).

20



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 12.00 (s, broad COOH, 1H); 10.24 (s, NH, 1H); 8.38 (d, J=5.8Hz, aromatic protons, 2H); 7.55 (d, J=5.8Hz, aromatic protons, 2H); 2.33 (t, J=7.2Hz, CH<sub>2</sub>CONHAr, 2H); 2.18 (t, J=7.2Hz, CH<sub>2</sub>COOH); 1.52 (m, 4H); 1.27 (m, 4H).

30




35


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200MHz), δ(ppm) 11.95 (s, COOH, 1H); 7.58 (d, J=8Hz); 3.50 (m, CH, 1H); 2.17 (t, J=7.2Hz, CH<sub>2</sub>COOH, 2H); 2.00 (t, J=7Hz, CH<sub>2</sub>CONH-, 2H); 1.60 (m, 4H); 1.46 (m, 6H); 1.20 (m, 8H).

40


In the same way the following compounds were prepared and characterized:




10 wherein  $n = 4, 5, 6, 7,$  and  $8$ ;  $R$  is hydrogen; 2-, 3-, and 4-cyano; 2-, 3-, and 4-nitro; 2-, 3-, and 4-methylcyano; 2-, 3-, and 4-trifluoromethyl; 2-, 3-, and 4-fluoro;



20 wherein  $n = 4, 5, 6, 7,$  and  $8$ ;




30 wherein  $n = 4, 5, 6, 7,$  and  $8$ ;




wherein  $n = 4, 5, 6, 7,$  and  $8$ ;

**SUBSTITUTE SHEET**


-36-



wherein  $n = 4, 5, 6, 7,$  and  $8;$




wherein  $n = 4, 5, 6, 7,$  and  $8;$



25 wherein R is 2-, 3-, and 4-carboxy; 2-, 3-, and 4-aminocarbonyl; 2-, 3-, and 4-methylaminocarbonyl; 2-, 3-, and 4-dimethylaminocarbonyl; 2-, 3-, and 4-chloro; 2-, 3-, and 4-bromo; 2-, 3-, and 4-iodo; 2-, 3-, and 4-methyl; 2-, 3-, and 4-methoxy; 2-, 3-, and 4-hydroxy; 2-, 3-, and 4-amino; and 2-, 3-, and 4-dimethylamino.

30

Compounds having the general structure:



**SUBSTITUTE SHEET**



wherein n = 4, 5, 6, and 7.

General Procedure A

5 A pyridine (500 mL) suspension of O-benzylhydroxylamine  
hydrochloride (3.2 g; 0.02 mol) and the corresponding  
diacid dichloride (0.04 mol) was stirred at room  
temperature for three days. Water (10 mL) was added and  
stirring was continued overnight. The solvent was  
10 evaporated and solid residue was purified by column  
chromatography on silica gel in tetrahydrofuran-methanol.  
The diacid product was dissolved in methanol (100 mL) and  
5% Pd-C (100 mg) was added. The reaction suspension was  
shaken overnight under hydrogen pressure (~50 psi). The  
15 catalyst was separated by filtration, solid residue was  
washed with hot methanol (5 x 50 ml). The combined  
methanolic filtrates were evaporated. The solid residue  
was slurried in acetone and filtered. The yield was 10-  
20%.

20


General procedure B

A pyridine (500 ml) solution of O-benzylhydroxylamine  
(2.46 g; 0.02 mol) and the corresponding dicarboxylic  
25 acid monobenzyl ester monoacid chloride (0.04 mol) was  
stirred at room temperature overnight. The solvent was  
evaporated. The semisolid residue was dissolved in  
chloroform (300 mL) and extracted with 5% hydrochloric  
acid (2 x 50 mL), 10% potassium hydroxide (3 x 100 mL),  
30 and water (2 x 100 mL). The organic layer was dried over  
anhydrous magnesium sulfate and evaporated. The solid  
residue was purified by column chromatography on silica  
gel in ethyl acetate. The tribenzyl product was  
dissolved in methanol (100 mL) and 5% Pd-C (100 mg) was  
35 added. The reaction suspension was shaken under hydrogen  
pressure (~50 psi) at room temperature overnight. The  
solid was separated by filtration and washed with hot

-38-

methanol (5 x 50 mL). The combined methanol filtrates were evaporated to solid residue. The solid residue was slurried in cooled acetone and filtered. The yield of target product was 30-60%.

5




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200MHz), δ(ppm) 11.53 (s, COOH, 1H); 2.41 (t, J=7.2Hz, CH<sub>2</sub>CON(OH)COCH<sub>2</sub>, 4H); 2.18 (t, J=7.0Hz, CH<sub>2</sub>COOH, 4H); 1.52 (m, 8h); 1.22 (m, H). MS (FAB, glycerin) 346(M + 1)

15

Compounds having the structure:

20



25


A pyridine (500 mL) solution of the monomethyl ester monoacid chloride of dicarboxylic acid (0.02 mol) and N,N'-dimethyl-1,ω-diaminoalkane (0.01 mol) was stirred at room temperature overnight. Solvent was evaporated and oily residue was dissolved in chloroform (300 mL). Chloroform solution was washed with water (3 x 50 mL), 10% potassium hydroxide (3 x 50 mL), 10% hydrochloric acid (3 x 50 mL), and again with water (3 x 50 mL). The organic layer was dried and evaporated. The oily residue was dissolved in potassium hydroxide (1.2 g; 0.021 mol) in 80% methanol (100 mL). The reaction mixture was refluxed two hours. The solvent was evaporated and solid residue was dissolved in water (50 mL) and extracted with chloroform (3 x 50 mL). Water solution was acidified to

30

35

**SUBSTITUTE SHEET**

pH~5 and concentrated (to volume of about 10 mL). The water solution or suspension was cooled down and precipitate was separated by filtration. The solid product was purified by crystallization from ethyl acetate. The yield was 40-60%.




$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 200 MHz),  $\delta$ (ppm) 8.15 (s, broad, COOH, 2H); 3.52 + 3.45 (2s,  $\text{CH}_2\text{N}$ , 4H); 3.01 + 2.93 (2s,  $\text{CH}_3\text{N}$ , 6H); 2.30 (4t,  $\text{CH}_2\text{CO}$ , 8H); 1.60 (m, 8H); 1.32 (m, 8H).

15  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 200 MHz),  $\delta$ (ppm) 3.44 + 3.336 + 3.36 (3s,  $\text{CH}_2\text{N}$ , 4H); 2.94 + 2.90 + 2.79 (3s,  $\text{CH}_3\text{N}$ , 6H); 2.27 + 2.23 + 2.12 (3t,  $\text{CH}_2\text{CO}$ , 8H); 1.46 (m, 8H); 1.23 (m, 8H).

Compounds having the structure:


20



25 A pyridine (500 mL) solution of 6-aminocaproic acid (2.6 g; 0.02 mol) and terephthaloyl chloride (2 g; 0.01 mol) was stirred at room temperature overnight (~12 hours), and at 90°C for 23 hours. The solvent was evaporated, and the solid residue was crystallized from water (10 mL)

30 four times. The yield was 800 mg (19%).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 200 MHz),  $\delta$ (ppm) 12.8 (s, broad, COOH, 2H); 8.54 + 7.72 (2t, NH, 2H); 3.24 + 2.98 (2m,  $\text{NHCH}_2$ , 4H); 2.20 + 2.03 (2m,  $\text{CH}_2\text{CO}$ , 4H); 1.50 (m, 8H); 1.32 (m, 4H).

35 Compound having the structure:




**SUBSTITUTE SHEET**

-40-

Into a mixture of aniline (2.75 g; 0.03 mol), hydroxylamine hydrochloride (2.08 g; 0.03 mol), and potassium hydroxide (5.50g; 0.09 mol) in 50% tetrahydrofuran (100 mL) was slowly added at room temperature a tetrahydrofurane (20 mL) solution of terephthaloyl chloride (6 g; 0.03 mol). The reaction suspension was stirred at room temperature for thirty minutes. The solvent was evaporated. The solid residue was slurried in hot methanol (1000 mL) and dried over anhydrous magnesium sulfate. The methanol solution was separated by filtration and filtrate was evaporated. The solid residue was slurried in 20 mL cooled methanol and filtered. The white crystals were washed with ether (5 x 50 mL) and dried. The yield was 4.6 g (39%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 11.35 (s, broad, NHOH, 1H); 10.35 (s, NHPH, 1H); 9.19 (s, NHOH, 1H); 8.03 (d, J=8Hz, terephthalic protons, 2H); 7.89 (d, J=8Hz, terephthalic protons, 2H); 7.82 (d, J=7.4Hz, ortho anilide protons, 2H); 7.34 (t, J=7.4Hz, meta anilide protons, 2H); 7.10 (t, J=7.4Hz, para anilide proton, 1H).

Compound having the structure:




A solution of 1,4-phenylenediacrylic acid (2.18 g; 0.01 mol) in thionyl chloride (50 mL; 81.55g; 0.68 mol) was refluxed overnight. The excess of thionyl chloride was evaporated. The solid was dissolved in tetrahydrofuran (20 mL), and added to a cooled (0°C) solution of potassium hydroxide (1.12 g; 0.02 mol) and aniline in 50% tetrahydrofuran. The reaction mixture was stirred at room temperature for thirty minutes. The solvent was evaporated. The solid residue was slurried in water and filtered. White crystals were dissolved in a small

**SUBSTITUTE SHEET**


amount of methanol and purified on a silica gel column in tetrahydrofuran. The yield was 315 mg (10%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 10.80 (s, NHOH, 1H); 10.23 (s, NHPH, 1H); 9.09 (s, NHOH, 1H); 7.69 (d, J=7.6Hz, ortho anilide protons, 2H); 7.64 (s, phenylene protons, 4H), 7.55 (d, J=15.8Hz, PhNHOCCH=CH-, 1H); 7.40 (d, J=15.8Hz, HONHOCCH=CH-, 1H); 7.33 (t, J=7.8Hz, meta anilide protons, 2H); 7.06 (t, J=7.2Hz, para anilide protons, 1H); 6.89 (d, J=15.8Hz, PhNHOCCH=CH-, 1H) 6.51 (d, J=15.8Hz, HONHOCCH=CH-, 1H).

Compounds having the structure:



wherein n = 4, 5, 6, 7, and 8.


20 A chloroform solution of triethylamine (1.4 mL; 1.0 g; 0.01 mol), the corresponding amine (0.01 mol) and diacid dichloride (0.005 mol) was stirred at room temperature for five hours. If the reaction mixture was clear, it was washed with water (5 x 100 mL). The organic layer was dried over anhydrous magnesium sulfate and evaporated to a solid residue. If in the course of reaction a precipitate was formed, the precipitate was separated by filtration. White crystals from filtration or solid residue from evaporation were crystallized from ethyl acetate, tetrahydrofuran, methanol, or their mixture. The yields were from 60-90%.



-42-

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.23 (s, NH, 2H); 7.82 (d,  $J=9\text{Hz}$ , aromatic protons, 4H), 7.60 (d,  $J=9\text{Hz}$ , aromatic protons, 4H), 2.31 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{CO}$ , 4H); 2.61 (m, 4H); 1.32 (m, 4H).


5



10

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.48 (s, NH, 2H); 8.18 (d,  $J=9.2\text{Hz}$ , aromatic protons, 4H); 7.81 (d,  $J=9.2\text{Hz}$ , aromatic protons, 4H); 2.37 (t,  $J=7.2\text{Hz}$ ,  $\text{CH}_2\text{CO}$ -, 4H); 1.60 (m, 4H); 1.33 (m, 4H).


15



20

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$  9.91 (s, NH, 2H), 7.58 (d,  $J=8.6\text{Hz}$ , aromatic protons, 4H); 7.26 (d,  $J=8.6\text{Hz}$ , aromatic protons, 4H); 3.94 (s,  $\text{CH}_2\text{CN}$ , 4H); 2.29 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{CO}$ -, 4H); 1.60 (m, 4H); 1.31 (m, 4H).


25



30

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.08 (s, CONHAr, 2H); 7.79 (d,  $J=8.6\text{Hz}$ , aromatic protons, 4H); 7.63 (d,  $J=8\text{Hz}$ , aromatic protons, 4H), 7.22 (s,  $\text{H}_3\text{CHNCO}$ -, 2H); 3.32 (s,  $\text{CH}_3$ , 6H); 2.31 (t,  $J=7\text{Hz}$ ,  $\text{CH}_2\text{C}$ -, 6H); 1.59 (m, 4H); 1.31 (m, 4H).

35




40


SUBSTITUTE SHEET



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 10.90 (s, broad, NHOH, 2H); 10.05 (s, NHAr, 2H); 8.90 (s, broad, NHOH, 2H); 7.68 (d, J=9Hz, aromatic protons, 4H); 7.62 (d, J=9Hz, aromatic protons, 4H); 2.31 (t, J=7.2Hz, CH<sub>2</sub>CO-, 4H); 1.59 (m, 4H); 1.30 (m, 4H).




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 10.06 (s, broad, NH, 2H); 8.71 (d, J=2.6Hz, aromatic protons, 2H); 7.31 (d + d, aromatic protons, 2H); 2.32 (t, J=7.4Hz, CH<sub>2</sub>CO-, 4H); 1.59 (m, 4H); 1.33 (m, 4H).



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 12.00 (s, broad, NH, 2H); 7.43 (d, J=3.6Hz, aromatic protons, 2H); 7.16 (d, J=3.6Hz, aromatic protons, 2H); 2.41 (t, J=7.2Hz, CH<sub>2</sub>CONH-, 4H); 1.58 (m, 4H); 1.28 (m, 4H).

In the similar manner, the following compounds were prepared and characterized:



wherein n = 4, 5, 6, 7, and 8;


40

all compounds are symmetrical wherein R is 2-, 3-, and 4-cyano; 2-, 3-, and 4-methylcyano; 2-, 3-, and 4-nitro,

-44-

2-, 3-, and 4-carboxy; 2-, 3-, and 4-aminocarbonyl; 2-, 3- and 4-methylaminocarbonyl; 2-, 3-, and 4-dimethylaminocarbonyl; and 2-, 3-, and 4-trifluoromethyl;


5



10


wherein R is 4-hydroxylaminocarbonyl; 4-methoxycarbonyl; 2-, 3-, and 4-chloro; 2-, 3-, and 4-fluoro; 2-, 3-, and 4-methyl; 2-, 3-, and 4-methoxy; 2,3-difluoro; 2,4-difluoro; 2,5-difluoro; 2,6-difluoro; 1,2,3-trifluoro, 3,4,5-trifluoro; 2,3,5,6-tetrafluoro; 2,3,4,5,6-pentafluoro.

15





20

25




30





Compounds having the structure:



25 wherein  $n = 4, 5, 6, 7, \text{ and } 8$ .

General procedure A

30 A diacid dichloride (0.01 mol) was added to a stirred solution of potassium hydroxide (1.68 g; 0.03 mol), hydroxylamine hydrochloride (0.7 g; 0.01 mol), and the corresponding aniline (0.01 mol) in 50% tetrahydrofuran (100 mL). The resulting reaction mixture was stirred at room temperature thirty minutes, and solvent was

35 evaporated to solid residue. The solid residue was slurried in methanol (~100 mL) and dried over anhydrous magnesium sulfate. The methanol solution was separated

by filtration and evaporated to a solid residue. The product was purified by column chromatography on silica gel in ethyl acetate-tetrahydrofuran (in most cases 3:1). The yields were 15-30%.

5

General procedure B

A solution of corresponding monomethyl ester of dicarboxylic acid (0.01 mol), oxaloyl chloride (0.03  
10 mol), and a few drops DMF in benzene (500 mL) was stirred at room temperature overnight. The solvent was evaporated and the oily residue was dissolved in dry benzene (3 x 50 mL) and evaporated again. The tetrahydrofuran (50 mL) solution of monoester monoacid  
15 chloride of the corresponding dicarboxylic acid was slowly added to a cooled solution of the corresponding amine (0.01 mol) and pyridine (1.6 mL; 1.6 g; 0.02 mol) in tetrahydrofuran (200 mL). The reaction mixture was stirred at room temperature for an hour. The solvent was  
20 evaporated, the residue was dissolved in chloroform (300 mL), and the chloroform solution was washed with 10% hydrochloric acid (3 x 50 mL), 10% potassium hydroxide (3 x 50 mL), and water (3 x 50 mL). The organic layer was dried over anhydrous magnesium sulfate and evaporated,  
25 yielding the pure monoester monoamide of dicarboxylic acid. The product was dissolved in 80% methanol with potassium hydroxide (0.56 g; 0.01 mol). The reaction mixture was refluxed two hours and evaporated to solid residue. The residue was dissolved in water (~20 mL) and  
30 acidified to ~pH 5 with 10% hydrochloric acid. The monoacid monoamide of the dicarboxylic acid was isolated by filtration of precipitate or extraction water solution with chloroform. The isolated monoacid monoamide of the dicarboxylic acid was mixed together with an equivalent  
35 amount of O-benzylhydroxylamine and 1,3-dicyclohexylcarbodiimide in pyridine (~100 mL per 0.01 mol of O-benzylhydroxylamine) and was stirred at room temperature

overnight. The solvent was evaporated and the solid residue was partitioned between chloroform (500 mL) and 10% hydrochloric acid (300 mL). The organic layer was washed with water (3 x 100 mL) and dried over anhydrous magnesium sulfate. The solvent was evaporated to solid residue. The solid residue was dissolved in large amounts of tetrahydrofuran and filtered through a short column of silica gel. The crude product was dissolved in methanol (100 mL) and 5% Pd-C was added. The reaction suspension was shaken under hydrogen pressure (~50 psi) overnight. The catalyst was separated by filtration and filtrate was evaporated to solid residue. The solid residue was slurried in hexanes and filtered. Mostly pure product was isolated in this way. If necessary further purification was achieved by column chromatography on silica gel with ethyl acetate-tetrahydrofuran. The yields were from 35% to 65%.


#### General procedure C

A pyridine (500 mL solution of O-benxylhydroxylamine (1.23; 0.01 mol), the corresponding amine (0.01 mol), and the dichloride of the dicarboxylic acid (0.01 mol) was stirred at room temperature overnight. The solvent was evaporated and the white solid residue contains, judged by <sup>1</sup>H NMR, two symmetrical amides and a target unsymmetrical one. The solid residue was slurried in methanol and dried over anhydrous magnesium sulfate. The filtrate was evaporated and the solid residue was dissolved in methanol (~100 mL). Into the methanol solution 5% Pd-C (100 mg) was added and the black suspension was shaken under hydrogen pressure (~50 psi) overnight. The catalyst was separated by filtration and the filtrate was evaporated. The product was isolated by column chromatography on silica with ethyl acetate-tetrahydrofuran. The yields were from 20% to 35%.

General procedure D

A chloroform solution of triethylamine (3 mL; 2.18 g; 0.0215 mol), the corresponding amine (0.01 mol),  
5 O-trimethylsilyl)hydroxylamine (1.05 g, 0.01 mol), and the corresponding diacid chloride of the dicarboxylic acid (0.01 mol) was stirred at room temperature overnight. The solvent was evaporated, the residue was dissolved in methanol (~10 mL), and into the methanol  
10 solution 10% ammonium chloride (~10 mL) was added. The resulting suspension was stirred at 50°C for two hours. The solvent was evaporated. The solid residue was slurried in methanol (300 mL) and dried over anhydrous magnesium sulfate. The methanol solution was separated  
15 by filtration and evaporated to a solid residue. The product was isolated by silica gel column chromatography with ethyl acetate-tetrahydrofuran. The yields were 20-33%.

20




25 Elemental analysis:      Calc.      63.62      7.63      10.60  
                                 Found      63.58      7.59      10.48

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 10.31 (s, NHOH, 1H); 9.83 (s, NHPH, 1H); 8.64 (s, NHOH, 1H); 7.57 (d, J=8.2Hz, ortho aromatic protons, 2H); 7.26 (t, J=8.4Hz, meta aromatic protons, 2H), 6.99 (t, J=7.4Hz, para aromatic protons, 1H); 2.27 (t, J=7.4Hz, CH<sub>2</sub>CONHPh, 2H); 1.93 (t, J=7.2Hz, CH<sub>2</sub>CONHOH, 2H); 1.52 (m, 4H); 1.26 (m, 4H). MS (Fab, Glycerin) 172, 204, 232, 249, 265, (100%, M + 1).



5




10

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.31 (s, NHOH, 1H); 10.08 (s, NHPH, 1H); 8.64 (s, NHOH, 1H); 7.78 (d,  $J=7.6\text{Hz}$ , aromatic protons, 1H); 7.66 (t,  $J=7.4\text{Hz}$ , aromatic protons, 1H); 7.48 (d,  $J=7.8\text{Hz}$ , aromatic protons, 1H); 7.29 (t,  $J=7.4\text{Hz}$ , aromatic protons, 1H); 2.34 (t,  $J=7\text{Hz}$ ,  $\text{CH}_2\text{CONHAr}$ , 2H); 1.93 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{CONHOH}$ , 2H); 1.58 (m, 4H); 1.27 (m, 4H).

15

20




25

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.31 (s, NHOH, 1H); 10.21 (s, NHPH, 1H); 8.65 (s, NHOH, 1H); 8.09 (s, aromatic proton, 1H); 7.77 (m, aromatic proton, 1H); 7.49 (m, aromatic proton, 1H); 2.31 (t,  $J=7.2\text{Hz}$ ,  $\text{CH}_2\text{CONHAr}$ , 2H); 1.93 (t,  $J=7.2\text{Hz}$ ,  $\text{CH}_2\text{CONHOH}$ , 2H); 1.51 (m, 4H).


30

35




40

$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.35 (s, NHAr, 1H); 10.31 (s, NHOH, 1H); 8.63 (s, NHOH + aromatic proton 2H); 7.88 (d,  $J=8\text{Hz}$ , aromatic protons, 2H); 7.57 (t,  $J=8\text{Hz}$ , aromatic proton, 1H); 2.33 (t,  $J=7.6\text{Hz}$ ,  $\text{CH}_2\text{CONHAr}$ , 2H); 1.93 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{CONHOH}$ , 2H), 1.52 (m, 4H); 1.27 (m, 4H).




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 10.33 (s, NHOH, 1H); 10.15 (s, NHAr, 1H); 10.09 (s, NHPh, 1H); 8.66 (s, NHOH, 1H); 7.91 (d, J=8.6Hz, aromatic protons, 2H); 7.76 (d, J=7.8Hz, ortho aniline protons, 2H); 7.71 (d, J=8.6Hz, aromatic protons, 2H); 7.33 (t, J=7.6Hz, meta anilide protons, 2H); 7.07 (t, J=7.4Hz, para anilide protons); 2.33 (t, J=7.5Hz, CH<sub>2</sub>NHAr, 2H); 1.93 (t, J=7.2Hz, CH<sub>2</sub>CNHH, 2H); 1.51 (m, 4H); 1.28 (m, 4H).



25  $^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 10.32 (s, NHOH, 1H);  
10.21 (s, NHAr, 1H); 8.65 (s, NHOH, 1H); 7.31 (d of d,  
J=10Hz(2.2Hz), aromatic protons, 2H); 6.84 (t of t,  
J=9.4Hz(2.4Hz), aromatic protons, 1H); 2.29 (t,  $\text{CH}_2\text{CONHAr}$ ,  
2H); 1.93 (t, J=7.2Hz,  $\text{CH}_2\text{CONHOH}$ , 2H); 1.51 (m, 4H); 1.26  
30 (m, 4H).


In the same manner the following compounds were prepared and characterized:





wherein n = 4, 5, 6, 7, and 8; and R is 2-, 3-, and 4-

-51-

cyano; 2-, 3-, and 4-methylcyano; 2-, 3-, and 4-nitro;  
 2-, 3-, and 4-carboxy; 2-, 3-, and 4-aminocarbonyl; 2-,  
 3-, and 4-methylaminocarbonyl; 2-, 3-, and 4-  
 dimethylaminocarbonyl; and 2-, 3-, and 4-  
 5 trifluoromethyl;




wherein R is 4-hydroxylaminocarbonyl; 4-methoxycarbonyl;  
 4-tetrazoyl; 2-, 3-, and 4-chloro; 2-, 3-, and 4-  
 15 fluoro; 2-, 3-, and 4-methyl; 2-, 3-, and 4-methoxy;  
 2,3-difluoro; 2,4-difluoro; 2,5-difluoro; 2,6-  
 difluoro; 1,2,3-trifluoro; 3,4,5-trifluoro; 2,4,5-  
 trifluoro; 2,4,6-trifluoro; 2,3,6-trifluoro; 2,3,5,6-  
 tetrafluoro; 2,3,4,5,6-pentafluoro; 2-, 3-, and 4-  
 20 phenyl; 2-, 3-, and 4-benzyloxy; 4-hexyl; and 4-t-  
 butyl;



SUBSTITUTE SHEET


Compounds having the structure:



10 wherein  $n = 4, 5, 6, 7,$  and  $8$ ; and  $R$  is hydrogen or methyl.


A diacid dichloride (0.01 mol) was added into a stirred solution of potassium hydroxide (1.68 g; 0.03 mol), aniline or N-methylaniline (0.01 mol), and dimethylamine hydrochloride (0.805 g; 0.01 mol) in 50% tetrahydrofuran (100 mL). The reaction mixture was stirred thirty minutes at room temperature. The solvent was partitioned between chloroform (400 mL) and water (300 mL). The organic layer was washed with 10% hydrochloric acid (3 x 100 mL), 10% potassium hydroxide (3 x 100 mL), and water (2 x 100 mL). The organic layer was dried over anhydrous magnesium sulfate and evaporated. The solid residue was slurried in hexanes and filtered. The yield were 25-34%.

25







$^1\text{H}$  NMR (DMSO- $d_6$ , 200 MHz),  $\delta$ (ppm) 9.82 (s, NHPH, 1H); 7.58 (d,  $J=7.6\text{Hz}$ , ortho aromatic protons, 2H); 7.26 (t,  $J=7.4\text{Hz}$ , meta aromatic protons, 2H); 6.99 (t,  $J=7.4\text{Hz}$ , para aromatic proton, 1H); 2.85 (d,  $J=28\text{Hz}$ ,  $\text{N}(\text{CH}_3)_2$ , 6H); 2.28 (t,  $J=7.2\text{Hz}$ ,  $\text{CH}_2\text{CO}$ , 2H); 2.24 (t,  $J=7.4\text{Hz}$ ,  $\text{CH}_2\text{CO}$ , 2H); 1.51 (m, 4H); 1.29 (m, 4H).

5



- 10 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 200 MHz), δ(ppm) 7.30 (m, C<sub>6</sub>H<sub>5</sub>, 5H); 3.13 (s, H<sub>3</sub>CNPh, 3H); 2.83 (d, J=26Hz, N(CH<sub>3</sub>)<sub>2</sub>, 6H); 2.17 (t, J=7.6Hz, CH<sub>2</sub>CON(CH<sub>3</sub>)<sub>2</sub>, 2H); 1.98 (t, J=7.4Hz, CH<sub>2</sub>CON(CH<sub>3</sub>)Ph, 2H); 1.41 (m, 4H); 1.11 (m, 4H).








TABLE 1

<u>CPD</u>	<u>Structure</u>	<u>Mol. Weight</u>	<u>Optimal Conc. (<math>\mu</math>M)</u>	<u>Benzidine Reactive Cells (%)</u>
				
1	n = 4 (known compound)	236	80	70
2	n = 5	250	20	84
3	n = 6	264	2.5	70
4	n = 7	278	20	8
5	n = 8	292	20	15
6		274	31	44
7		274	31	52
8		294	12.5	32

SUBSTITUTE SHEET












TABLE 1 (continued)

<u>CPD</u>	<u>Structure</u>	<u>Mol. Weight</u>	<u>Optimal Conc. (<math>\mu</math>M)</u>	<u>Benzidine Reactive Cells (%)</u>
9		225	50	20
10		355	250	26
11		216	60	53
12		189	250	35
13		203	60	17
14		156	125	30
15		218	20	43

SUBSTITUTE SHEET

TABLE 1 (continued)

<u>CPD</u>	<u>Structure</u>	<u>Mol. Weight</u>	<u>Optimal Conc. (<math>\mu</math>M)</u>	<u>Benzidine Reactive Cells (%)</u>
16		270	8	35
17		256	62	30
18		260	31	38
19		278	5	24
20	 R = 4-methyl	273	20	52
21	 R = 4-cyano	289	7	70
22	 R = 3-cyano	289	5	55
23	 R = 2-cyano	289	16	65
24	 R = 3-nitro	309	5	30

SUBSTITUTE SHEET

TABLE 1 (continued)

<u>CPD</u>	<u>Structure</u>	<u>Mol. Weight</u>	<u>Optimal Conc. (<math>\mu</math>M)</u>	<u>Benzidine Reactive Cells (%)</u>
25	R = 4-nitro	309	0.8	30
26	R = 3-trifluoromethyl	332	30	30
27	R = 4-trifluoromethyl	332	5	47
28	R = 2-amino	279	20	54
29	R = 4-cyanomethyl	303	1	30
30	R = 3-chloro	298.5	2	33
31	R = 4-azido ( $N_3$ )	304	2	47
32	R = 2-fluoro	282	4	65
33	R = 3-fluoro	282	1	25
34	R = 4-fluoro	282	4	43
35	R = 4-benzyloxy	370	4	20
36	R = 4-methoxycarbonyl	322	4	28
37	R = 4-methylaminocarbonyl	321	30	16
38	R = 2-bromo	343	8	45
39	R = 2-chloro	298.5	4	34
40	R = 4-bromo	343	1.6	47

SUBSTITUTE SHEET

TABLE 1 (continued)

<u>CPD</u>	<u>Structure</u>	<u>Mol. Weight</u>	<u>Optimal Conc. (<math>\mu</math>M)</u>	<u>Benzidine Reactive Cells (%)</u>
41	R = 2,3-difluoro	300	8	24
42	R = 2,4,5-trifluoro	318	8	36
43	R = 2,3,6-trifluoro	318	31	53
44	R = 2,4,6-trifluoro	318	16	47
45	R = 2,4-difluoro	300	6	60
46	R = 2,3,4,5,6-pentafluoro	354	31	53
47	R = 3,4-difluoro	300	4	61
48	R = 3,4,5-trifluoro	318	8	55
49	R = 2,5-difluoro	300	4	70
50	R = 3,5-difluoro	300	2	73
51	R = 2-methoxy	294	8	36
52	R = 3-methoxy	294	6	38
53	R = 4-methoxy	294	6	37






TABLE 1 (continued)

CPD	Structure	Mol. Weight	Optimal Conc. ( $\mu$ M)	Benzidine Reactive Cells (%)
55		256	30	53
				
56	R = 4-trifluoromethyl	460	50	20
57	R = 4(N)-hydroxylamino-carbonyl	442	8	10
58	R = 4-cyanomethyl	402	50	25
59	R = 2,4-difluoro	396	500	54
60	R = 2,6-difluoro	396	100	21
61	R = 3,5-difluoro	396	125	31
62	R = 2,3,6-trifluoro	432	250	28
63	R = 2,4,6-trifluoro	432	125	35
64	R = 2,3,4,5,6-pentafluoro	504	125	13
65	R = 4-nitro	414	25	14

SUBSTITUTE SHEET

TABLE 1 (continued)

CPD	Structure	Mol. Weight	Optimal Conc. ( $\mu$ M)	Benzidine Reactive Cells (%)
66	$\begin{array}{c} \text{O} \quad \text{CH}_3 \quad \text{CH}_3 \quad \text{O} \\ \parallel \quad   \quad   \quad \parallel \\ (\text{H}_3\text{C})_2\text{N}-\text{C}-\text{CH}-(\text{CH}_2)_5-\text{CH}-\text{C}-\text{N}(\text{CH}_3)_2 \end{array}$	270	1250	80
67	$\begin{array}{c} \text{O} \quad \text{CH}_3 \quad \text{CH}_3 \quad \text{O} \\ \parallel \quad   \quad   \quad \parallel \\ (\text{H}_3\text{C})_2\text{N}-\text{C}-\text{CH}-(\text{CH}_2)_4-\text{CH}-\text{C}-\text{N}(\text{CH}_3)_2 \end{array}$	256	2500	90
68	$\begin{array}{c} \text{O} \quad \text{CH}_3 \quad \text{O} \\ \parallel \quad   \quad \parallel \\ \text{HOHN}-\text{C}-(\text{CH}_2)_2-\text{CH}-(\text{CH}_2)_2-\text{C}-\text{NHOH} \end{array}$	204	125	56
69	$\begin{array}{c} \text{O} \quad \text{CONHOH} \quad \text{O} \\ \parallel \quad   \quad \parallel \\ \text{HOHN}-\text{C}-(\text{CH}_2)_5-\text{CH}-(\text{CH}_2)_5-\text{C}-\text{NHOH} \end{array}$	333	60	40
70	$\begin{array}{c} \text{O} \quad \text{F} \quad \text{O} \\ \parallel \quad   \quad \parallel \\ \text{HOHN}-\text{C}-(\text{CH}_2)_2-\text{CH}-(\text{CH}_2)_2-\text{C}-\text{NHOH} \\ \quad \quad \quad   \\ \quad \quad \quad \text{F} \end{array}$	226	160	19



TABLE 1 (continued)


<u>CPD</u>	<u>Structure</u>	<u>Mol. Weight</u>	<u>Optimal Conc. (<math>\mu</math>M)</u>	<u>Benzidine Reactive Cells (%)</u>
				
71	n = 4	310	100	8
72	n = 5	324	250	10
73	n = 6	338	50	7
74	n = 7	352	100	10
75	n = 8	366	100	10

TABLE 2

Induction of Differentiation of HL-60

<u>CPD</u>	Mol. <u>Weight</u>	Optimal <u>Conc. (<math>\mu</math>M)</u>	NBT <u>Positive (%)</u>
2	250	7	22
3	264	1	21
6	274	20	30
7	274	20	21
22	289	1.7	28
21	289	2	6
26	332	6	27
25	309	3	18
36	322	1	32
31	304	2.5	7
29	303	1	15
43	318	2	20

**References:**

1. Sporn, M.B., Roberts, A.B., and Driscoll, J.S. (1985) in Cancer: Principles and Practice of Oncology, eds. Hellman, S., Rosenberg, S.A., and DeVita, V.T., Jr., Ed. 2, (J.B. Lippincott, Philadelphia), P. 49.  
5
2. Breitman, T.R., Selonick, S.E., and Collins, S.J. (1980) Proc. Natl. Acad. Sci. USA 77: 2936-2940.  
10
3. Olsson, I.L. and Breitman, T.R. (1982) Cancer Res. 42: 3924-3927.
4. Schwartz, E.L. and Sartorelli, A.C. (1982) Cancer Res. 42: 2651-2655.  
15
5. Marks, P.A., Sheffery, M., and Rifkind, R.A. (1987) Cancer Res. 47: 659.  
20
6. Sachs, L. (1978) Nature (Lond.) 274: 535.
7. Friend, C., Scher, W., Holland, J.W., and Sato, T. (1971) Proc. Natl. Acad. Sci. (USA) 68: 378-382.  
25
8. Tanaka, M., Levy, J., Terada, M., Breslow, R., Rifkind, R.A., and Marks, P.A. (1975) Proc. Natl. Acad. Sci. (USA) 72: 1003-1006.
9. Reuben, R.C., Wife, R.L., Breslow, R., Rifkind, R.A., and Marks, P.A. (1976) Proc. Natl. Acad. Sci. (USA) 73: 862-866.  
30
10. Abe, E., Miyaura, C., Sakagami, H., Takeda, M., Konno, K., Yamazaki, T., Yoshika, S., and Suda, T. (1981) Proc. Natl. Acad. Sci. (USA) 78: 4990-4994.  
35

11. Schwartz, E.L., Snoddy, J.R., Kreutter, D., Rasmussen, H., and Sartorelli, A.C. (1983) Proc. Am. Assoc. Cancer Res. 24: 18.
- 5 12. Tanenaga, K., Hozumi, M., and Sakagami, Y. (1980) Cancer Res. 40: 914-919.
13. Lotem, J. and Sachs, L. (1975) Int. J. Cancer 15: 731-740.
- 10 14. Metcalf, D. (1985) Science, 229: 16-22.
15. Scher, W., Scher, B.M., and Waxman, S. (1983) Exp. Hematol. 11: 490-498.
- 15 16. Scher, W., Scher, B.M., and Waxman, S. (1982) Biochem. & Biophys. Res. Comm. 109: 348-354.
17. Huberman, E. and Callahan, M.F. (1979) Proc. Natl. Acad. Sci. (USA) 76: 1293-1297.
- 20 18. Lottem, J. and Sachs, L. (1979) Proc. Natl. Acad. Sci. (USA) 76: 5158-5162.
- 25 19. Terada, M., Epner, E., Nudel, U., Salmon, J., Fibach, E., Rifkind, R.A., and Marks, P.A. (1978) Proc. Natl. Acad. Sci. (USA) 75: 2795-2799.
20. Morin, M.J. and Sartorelli, A.C. (1984) Cancer Res. 44: 2807-2812.
- 30 21. Schwartz, E.L., Brown, B.J., Nierenberg, M., Marsh, J.C., and Sartorelli, A.C. (1983) Cancer Res. 43: 2725-2730.
- 35 22. Sugano, H., Furusawa, M., Kawaguchi, T., and Ikawa, Y. (1973) Bibl. Hematol. 39: 943-954.

23. Ebert, P.S., Wars, I., and Buell, D.N. (1976) Cancer Res. 36: 1809-1813.
- 5 24. Hayashi, M., Okabe, J., and Hozumi, M. (1979) Gann 70: 235-238.
25. Fibach, E., Reuben, R.C., Rifkind, R.A., and Marks, P.A. (1977) Cancer Res. 37: 440-444.
- 10 26. Melloni, E., Pontremoli, S., Damiani, G., Viotti, P., Weich, N., Rifkind, R.A., and Marks, P.A. (1988) Proc. Natl. Acad. Sci. (USA) 85: 3835-3839.
- 15 27. Reuben, R., Khanna, P.L., Gazitt, Y., Breslow, R., Rifkind, R.A., and Marks, P.A. (1978) J. Biol. Chem. 253: 4214-4218.
- 20 28. Marks, P.A. and Rifkind, R.A. (1988) International Journal of Cell Cloning 6: 230-240.
- 25 29. Melloni, E., Pontremoli, S., Michetti, M., Sacco, O., Cakiroglu, A.G., Jackson, J.F., Rifkind, R.A., and Marks, P.A. (1987) Proc. Natl. Acad. Sciences (USA) 84: 5282-5286.
- 30 30. Marks, P.A. and Rifkind, R.A. (1984) Cancer 54: 2766-2769.
- 30 31. Egorin, M.J., Sigman, L.M. VanEcho, D.A., Forrest, A., Whitacre, M.Y., and Aisner, J. (1987) Cancer Res. 47: 617-623.
- 35 32. Rowinsky, E.W., Ettinger, D.S., Grochow, L.B., Brundrett, R.B., Cates, A.E., and Donehower, R.C. (1986) J. Clin. Oncol. 4: 1835-1844.


-66-

33. Rowinsky, E.L. Ettinger, D.S., McGuire, W.P., Noe, D.A., Grochow, L.B., and Donehower, R.C. (1987) Cancer Res. 47: 5788-5795.
- 5 34. Callery, P.S., Egorin, M.J., Geelhaar, L.A., and Nayer, M.S.B. (1986) Cancer Res. 46: 4900-4903.
- 10 35. Young, C.W. Fanucchi, M.P., Walsh, T.B., Blatzer, L., Yaldaie, S., Stevens, Y.W., Gordon, C., Tong, W., Rifkind, R.A., and Marks, P.A. (1988) Cancer Res. 48: 7304-7309.
- 15 36. Andreeff, M., Young, C., Clarkson, B., Fetten, J., Rifkind, R.A., and Marks, P.A. (1988) Blood 72: 186a.
- 20 37. Marks, P.A., Breslow, R., Rifkind, R.A., Ngo, L., and Singh, R. (1989) Proc. Natl. Acad. Sci. (USA) 86: 6358-6362.
38. Breslow, R., Jursic, B., Yan, Z.F., Friedman, E., Leng, L., Ngo, L., Rifkind, R.A., and Marks, P.A. (1991) Proc. Natl. Acad. Sci. (USA) 88: 5542-5546.
- 25 39. Ohta, Y., Tanaka, M., Terada, M., Miller, O.J., Bank, A., Marks, P.A., and Rifkind, R.A. (1976) Proc. Natl. Acad. Sci. (USA) 73: 1232-1236.
- 30 40. Collins, S.J., Gallo, R.C., and Gallagher, R.E. (1978) Nature (London) 270; 405-409.
41. Synder, S.W., Egorin, M.J., Geelhaar, L.A., Hamburger, A.W., and Callery, P.S. (1988) Cancer Res. 48; 3613-3616.



What is claimed is:


1. A compound having the structure:



10 wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other; when  $R_1$  and  $R_2$  are the same, each is a substituted or unsubstituted arylamino, cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; when  $R_1$  and  $R_2$  are different,  $R_1 = R_3\text{-N-}R_4$ ,  
 15 wherein each of  $R_3$  and  $R_4$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy,  
 20 arylalkyloxy, or pyridine group, or  $R_3$  and  $R_4$  bond together to form a piperidine group and  $R_2$  is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and  $n$  is an integer from about 4 to about 8.

25

2. A compound of claim 1 having the structure:




35 wherein each of  $R_3$  and  $R_4$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy,

-68-

- arylalkyloxy, or pyridine group, or  $R_3$  and  $R_4$  bond together to form a piperidine group;  $R_2$  is a hydroxylamino, hydroxyl, amino, alkylamino, dialkylamino or alkyloxy group; and  $n$  is an integer from about 4 to about 8.
- 5
3. A compound of claim 2, wherein  $R_2$  is a hydroxylamino, hydroxyl, amino, methylamino, dimethylamino, or methoxy group and  $n$  is 6.
- 10
4. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a substituted or unsubstituted phenyl group.
- 15
5. A compound of claim 4, wherein the phenyl group is substituted with a methyl, cyano, nitro, trifluoromethyl, amino, aminocarbonyl, methylcyano, chloro, fluoro, bromo, iodo, 2,3-difluoro, 2,4-difluoro, 2,5-difluoro, 3,4-difluoro, 3,5-difluoro, 2,6-difluoro, 1,2,3-trifluoro, 2,3,6-trifluoro, 2,4,6-trifluoro, 3,4,5-trifluoro, 2,3,5,6-tetrafluoro, 2,3,4,5,6-pentafluoro, azido, hexyl, t-butyl, phenyl, carboxyl, hydroxyl, methoxy, phenyloxy, benzyloxy, phenylaminooxy, phenylaminocarbonyl, methoxycarbonyl, methylaminocarbonyl, dimethylamino, dimethylamino-carbonyl, or hydroxylaminocarbonyl group.
- 20
- 25
- 30
6. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a cyclohexyl group.
7. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a methoxy group.
- 35
8. A compound of claim 3, wherein  $R_3$  and  $R_4$  bond together to form a piperidine group.

SUBSTITUTE SHEET

9. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a hydroxyl group.
- 5 10. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a benzyloxy group.
11. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a  $\delta$ -pyridine group.
- 10 12. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a  $\beta$ -pyridine group.
13. A compound of claim 3, wherein  $R_4$  is a hydrogen atom and  $R_3$  is a  $\alpha$ -pyridine group.
- 15 14. A compound of claim 3, wherein  $R_3$  and  $R_4$  are both methyl groups.
15. A compound of claim 3, wherein  $R_4$  is a methyl group and  $R_3$  is a phenyl group.
- 20 16. A compound of claim 1 having the structure:



30 wherein R is a substituted or unsubstituted arylamino, cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiozoleamino group; and n is an integer from about 4 to about 8.


- 35 17. A compound of claim 16, wherein R is a substituted or unsubstituted phenylamino group.

18. A compound of claim 17, wherein the phenylamino

group is substituted with a cyano, methylcyano, nitro, carboxyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, trifluoromethyl, hydroxylaminocarbonyl, N-hydroxylaminocarbonyl, methoxycarbonyl, chloro, fluoro, methyl, methoxy, 2,3-difluoro, 2,4-difluoro, 2,5-difluoro, 2,6-difluoro, 3,5-difluoro, 2,3,6-trifluoro, 2,4,6-trifluoro, 1,2,3-trifluoro, 3,4,5-trifluoro, 2,3,4,5-tetrafluoro, or 2,3,4,5,6-pentafluoro group.

19. A compound of claim 16, wherein R is a cyclohexylamino group.


20. A compound having the structure:



wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; R is a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

21. A compound of claim 20, wherein each of X, Y, and R is a hydroxyl group and each of m and n is 5.

22. A compound having the structure:




wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m, n, and o are independently the same as or different from each other and are each an integer from about 0 to about 8.

23. A compound of claim 22, wherein each of X and Y is a hydroxyl group and each of R<sub>1</sub> and R<sub>2</sub> is a methyl group.

24. A compound of claim 23, wherein each of n and o is 6, and m is 2.


25. A compound having the structure:



wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino,

arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of  $m$  and  $n$  are independently the same as or different from each other and are each an integer from about 0 to about 8.

26. A compound having the structure:




15

wherein each of  $X$  and  $Y$  are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; and each of  $m$  and  $n$  are independently the same as or different from each other and are each an integer from about 0 to about 8.

27. A compound of claim 26, wherein each of  $X$  and  $Y$  is a hydroxyl group and each of  $m$  and  $n$  is 5.

28. A compound having the structure:




35



wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, or aryloxy group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

29. A compound having the structure:




wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; and n is an integer from about 0 to about 8.

30. A compound of claim 29, wherein each of X and Y is a dimethylamino group and n is 5.

31. A compound of claim 29, wherein each of X and Y is a dimethylamino group and n is 4.

32. A compound having the structure:



10 wherein each of X and Y are independently the same as or different from each other and are a hydroxyl, amino or hydroxylamino group, a substituted or unsubstituted alkyloxy, alkylamino, dialkylamino, arylamino, alkylarylamino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group; each of R<sub>1</sub> and R<sub>2</sub> are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted alkyl, aryl, alkyloxy, aryloxy, carbonylhydroxylamino, or fluoro group; and each of m and n are independently the same as or different from each other and are each an integer from about 0 to about 8.

15

20

33. A compound of claim 32, wherein each of X and Y is a hydroxylamino group; R<sub>1</sub> is a methyl group; R<sub>2</sub> is a hydrogen atom; and each of m and n is 2.


25

34. A compound of claim 32, wherein each of X and Y is a hydroxylamino group; R<sub>1</sub> is a carbonylhydroxylamino group; R<sub>2</sub> is a hydrogen atom; and each of m and n is 5.

30

35. A compound of claim 32, wherein each of X and Y is a hydroxylamino group; each of R<sub>1</sub> and R<sub>2</sub> is a fluoro group; and each of m and n is 2.

35 36. A compound having the structure:




**SUBSTITUTE SHEET**

wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

37. A compound of claim 36, wherein  $R_1$  is a phenylamino group and  $R_2$  is a hydroxylamino group.


38. A compound having the structure:



wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

39. A compound of claim 38, wherein  $R_1$  is phenylamino group and  $R_2$  is hydroxylamino group.

40. A compound having the structure:



wherein each of  $R_1$  and  $R_2$  are independently the same as or different from each other and are a hydroxyl, alkyloxy, amino, hydroxylamino, alkylamino, dialkylamino, arylamino, alkylaryl amino, alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

alkyloxyamino, aryloxyamino, alkyloxyalkylamino, or aryloxyalkylamino group.

- 5 41. A compound of claim 40, wherein  $R_1$  is a hydroxylamino group.
42. A compound of claim 40, wherein  $R_2$  is a hydroxylamino group.
- 10 43. A method of selectively inducing terminal differentiation of neoplastic cells and thereby inhibiting proliferation of such cells which comprises contacting the cells under suitable conditions with an effective amount of the compound of claim 1, 2, 16, 20, 22, 25, 26, 28, 29, 32, 36, 15 38, or 40, effective to selectively induce terminal differentiation.
- 20 44. A method of treating a patient having a tumor characterized by proliferation of neoplastic cells which comprises administering to the patient an effective amount of the compound of claim 1, 2, 16, 20, 22, 25, 26, 28, 29, 32, 36, 38, or 40, effective to selectively induce terminal differentiation of 25 such neoplastic cells and thereby inhibit their proliferation.
- 30 45. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compound of claim 1, 2, 16, 20, 22, 25, 26, 28, 29, 32, 36, 38, or 40.
- 35 46. A pharmaceutical composition of claim 45, wherein the effective amount is an amount effective to selectively induce terminal differentiation of suitable neoplastic cells and less than an amount

which causes toxicity in a patient.

47. A pharmaceutical composition of claim 45 in combination with an antitumor agent.

5

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US92/08454

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : Please See Extra Sheet.  
US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A 2,895,991 (RANDALL ET AL) 21 JULY 959, COL. 1 LINES 55 +	1-19,29-35,43-47
Y	US,A 4,056,524 (WALKER) 1 NOVEMBER 1977, COL. 1 LINES 1+5	1-19,29-35,43-47
Y	US, A, 4,480,125 (HAAS ET AL) 30 OCTOBER 1984, COL. 2 LINES 15 +	1-19,29-35,43-47
Y	US, A, 4,537,781 (DARLING) 27 AUGUST 1985, COL. 2 LINES 3 +	1-19,29-35,43-47

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

<p>* Special categories of cited documents:</p>		<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p>	
"A"	document defining the general state of the art which is not considered to be part of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

11 January 1993

Date of mailing of the international search report

11 FEB 1993

Name and mailing address of the ISA/  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

*Raymond Covington*  
RAYMOND COVINGTON

Telephone No. (703) 308-4704



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US92/08454

**A. CLASSIFICATION OF SUBJECT MATTER:**  
IPC (5):

CO7D 473/02 211/26 211/28 211/30 277/20 277/44 277/02; CO7C 271/02 271/08 271/10 271/12 271/14 271/18  
229/02 229/24 229/26 229/34 229/40 233/04 233/05 233/06 233/07 233/12 233/31 233/47 233/51 235/74 235/76  
237/06 237/08 239/00

**A. CLASSIFICATION OF SUBJECT MATTER:**  
US CL :

544/277;546/190,224,225;548/195;560/29,30,31,55,104,105,159,160,170,190,192,205,215,;562/433,450,465,470,555,  
582,583,587,595,596,;564/152,155,158,160,161,169,170,171,182,188,192,193,194,197,199,200,201,202,204,209

**B. FIELDS SEARCHED**

Minimum documentation searched

Classification System: U.S.

544/277;546/190,224,225;548/195;560/29,30,31,55,104,105,159,160,170,190,192,205,215,;562/433,450,465,470,555,  
582,583,587,595,596,;564/152,155,158,160,161,169,170,171,182,188,192,193,194,197,199,200,201,202,204,209

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING**

This ISA found multiple inventions as follows:

- I. CLAIMS 1-19 AND 29-35, DRAWN TO RING CONTAINING BIS-AMINO SUBSTITUTED ALKYL DERIVATIVES.
- II. CLAIMS 2 AND 20-21, DRAWN TO CARBOXYLIC ACID AMIDE SUBSTITUTED DERIVATIVES.
- III. CLAIM 22-24 DRAWN TO CARBAZIDE DERIVATIVES.
- IV. CLAIM 25 DRAWN TO BIS-CARBONYL CONTAINING AMIDE ALKYL DERIVATIVES.
- V. CLAIMS 26 AND 27 DRAWN TO BIS-AMIDE SUBSTITUTED ARYL DERIVATIVES.
- VI. CLAIM 28 DRAWN TO BIS-CARBONYLAMINE SUBSTITUTED ARYL DERIVATIVES.
- VII. CLAIM 36-37 DRAWN TO DICARBOXYLIC ACID ARYL DERIVATIVES.
- VIII. CLAIMS 38 AND 39 DRAWN TO VINYL-CARBONYL ARYL DERIVATIVES.
- IX. CLAIMS 40-42 DRAWN TO CARBOXYLIC ACID VINYL DERIVATIVES.